

## Tylenol

**\*NO Tylenol for infants under 3 months old unless discussed with doctor first.**

### Infant Concentrated Drops (80 mg per 0.8 ml)

|             |        |                 |
|-------------|--------|-----------------|
| 8 - 11 lbs  | 0.6 ml | every 4-6 hours |
| 12 - 14 lbs | 0.8 ml | every 4-6 hours |
| 15 - 17 lbs | 1 ml   | every 4-6 hours |
| 18 - 21 lbs | 1.2 ml | every 4-6 hours |
| 22 - 26 lbs | 1.6 ml | every 4-6 hours |
| 27 - 30 lbs | 2 ml   | every 4-6 hours |

### Children's Suspension (160 mg per 5 ml)

|             |      |                 |
|-------------|------|-----------------|
| 20 - 24 lbs | 4 ml | every 4-6 hours |
| 25 - 30 lbs | 5 ml | every 4-6 hours |
| 31 - 35 lbs | 7 ml | every 4-6 hours |
| 36 - 40 lbs | 8 ml | every 4-6 hours |
| 41 - 45 lbs | 9 ml | every 4-6 hours |
| 46 - 50 lbs | 10ml | every 4-6 hours |

## Benadryl

**\*NONE under 6 months old**

### Benadryl Elixir (12.5 mg per 5 ml)

|             |      |                 |
|-------------|------|-----------------|
| 16 - 20 lbs | 3 ml | every 6-8 hours |
| 21 - 25 lbs | 4 ml | every 6-8 hours |
| 26 - 32 lbs | 5 ml | every 6-8 hours |
| 33 - 40 lbs | 7 ml | every 6-8 hours |
| 41 - 50 lbs | 8 ml | every 6-8 hours |